arXiv:math-ph/0103003v2 6 Apr 2001

A Remark on Topological Charges over the Fuzzy Sphere

Harald Grosse* and Christian W. Rupp†

Institut für Theoretische Physik, Universität Wien,
Boltzmanng. 5, A–1090 Wien, Austria,
Institut für Theoretische Physik, Universität Leipzig
Augustusplatz 10–11, D–04109 Leipzig, Germany

Abstract

We determine the Chern characters of two projective modules over the fuzzy sphere and calculate the corresponding topological charges (Chern numbers). These turn out to have corrections—compared to the commutative limit—induced by the noncommutative structure of the three coordinates.

1 Introduction and Overview

Classical gauge field theories exhibit interesting features connected with the geometry and topology of nontrivial fiber bundles (over space or space–time). Examples are monopole and instanton solutions. In noncommutative geometry the situation is similar since the algebraic aspect of nontriviality is nothing but the projectivity of certain algebra modules.

For the fuzzy sphere this case has first been analyzed by one of us (HG) et. al. in [1] (see also [2]), leading to scalar and spinor field configurations in monopole backgrounds. A different approach using spectral triples and their Dirac operators has been used in [3] and [4]. Recently [5] there have been attempts to calculate Chern numbers applying a noncommutative version of the projectors for the 2–sphere used in [6] and [7]. In our present analysis we use this setup to show that the Chern characters obtained in this way

---

*E-mail: grosse@thp.univie.ac.at

†E-mail: cwrupp@itp.uni-leipzig.de

give rise to non-integer (fuzzy) topological charges, becoming integer in the commutative limit.

In section 2 we briefly review the Chern character on projective modules over algebras equipped with a differential calculus, primarily to settle notation. Further information can be found e.g. in [8], [9] and [10]. Then in section 3 we review the complex line bundles over the 2–sphere, and in section 4 we generalize the bundles with topological charge ±1 to the fuzzy sphere, resulting in fuzzy line bundles, their Chern characters and Chern numbers.

2 The Chern Character

Let $\mathcal{A}$ be a complex unital not necessarily commutative $C^\star$-algebra and denote $\mathcal{A}\otimes\mathbb{C}^n$ by $\mathcal{A}^n$. Then any projector (selfadjoint idempotent) $p \in M_n(\mathcal{A})$, the $n\times n$-matrices with coefficients in $\mathcal{A}$, defines a (finitely generated) projective right $\mathcal{A}$-module $E = p\mathcal{A}^n$. Elements ψ of E can be viewed as $\psi \in \mathcal{A}^n$ with $p\psi = \psi$. If $\mathcal{A}$ is further endowed with a differential calculus $(\Omega^*(\mathcal{A}), d)$, the Grassmann connection $\nabla : E \to E\otimes_{\mathcal{A}}\Omega^1(\mathcal{A})$ of E is defined by $\nabla = p\circ d$. It satisfies $\nabla(\psi f) = (\nabla\psi)f + \psi\otimes_{\mathcal{A}} df$ for all $f \in \mathcal{A}$, $\psi \in E$. After extending ∇ to $E\otimes_{\mathcal{A}}\Omega^1(\mathcal{A})$ one can define the $\mathcal{A}$-linear map

$$\nabla^2 : E \to E\otimes_{\mathcal{A}}\Omega^2(\mathcal{A}),$$

called the curvature of ∇. Evaluating ∇^2 one finds $\nabla^2 = p(dp)(dp)$. The Chern character of E is the set of

$$\mathrm{Ch}_q(p) := \frac{1}{q!}\mathrm{Tr}\, p(dp)^{2q}, \tag{1}$$

which are a cocycles and provide equivalence classes in $H^{2q}(\mathcal{A})$. $\mathrm{Ch}_0(p) = \mathrm{Tr}\, p$ simply gives the rank of the module. In the case of the 2–sphere and the fuzzy sphere the highest non-vanishing component of the Chern character is Ch_1. If $\mathcal{A}$ is a commutative algebra and E_1 and E_2 are both projective $\mathcal{A}$–modules, then the first component of the Chern character on the module tensor product has the property

$$\mathrm{Ch}_1(E_1\otimes_{\mathcal{A}} E_2) = \sum_{i+j=1} \mathrm{Ch}_i(E_1)\mathrm{Ch}_j(E_2), \tag{2}$$

as it is the case for bundles of higher topological charge over the 2–sphere.

3 Line Bundles over the 2–Sphere

In this section we give an instant derivation of all inequivalent complex line bundles over S^2. Define the 2-sphere by

$$(x_1, x_2, x_3) \in \mathbb{R}^3 \quad \text{with} \quad (x_1)^2 + (x_2)^2 + (x_3)^2 = 1$$

and denote by $\mathcal{A} = C^\infty(S^2)$ the commutative $\mathbb{C}$-algebra of smooth functions on it. The Bott projector $p = p^2 = p^\dagger$ is

$$p = \frac{1}{2}(1 + \sigma_a x_a) \in M_2(\mathcal{A}), \tag{3}$$

a 2×2-matrix with coefficients in $\mathcal{A}$, where σ_a are the three Pauli matrices. p defines a line bundle—note that $\operatorname{Tr} p = 1$—over S^2 through the right $\mathcal{A}$-module of smooth sections $E = p\mathcal{A}^2$. The first component $F \in \Omega^2(S^2)$ of the Chern character of E is given by

$$\begin{aligned} F &= \operatorname{Tr} p(dp)(dp) = \frac{1}{8}\operatorname{Tr}(1 + \sigma_a x_a)(\sigma_b dx_b)(\sigma_c dx_c) \\ &= \frac{1}{8}\operatorname{Tr}(\sigma_b\sigma_c + \sigma_a x_a \sigma_b \sigma_c) dx_b \wedge dx_c. \end{aligned}$$

Using $\sigma_b\sigma_c = \delta_{bc} + i\epsilon_{bcd}\sigma_d$ and the tracelessness of the Pauli matrices this expression can be seen to be equal to

$$F = \frac{1}{4} i\epsilon_{abc} x_a dx_b \wedge dx_c. \tag{4}$$

Since $\frac{1}{2}\epsilon_{abc}x_a dx_b \wedge dx_c = \sin\theta d\theta \wedge d\phi$ it holds that

$$\int_{S^2} \frac{1}{8\pi}\epsilon_{abc} x_a dx_b \wedge dx_c = 1,$$

and therefore the first Chern number (topological charge) $c_1(p)$ of the bundle E is found to be

$$c_1(p) = \frac{1}{2\pi i}\int_{S^2} F = 1. \tag{5}$$

Notice that $c_1(p^{\mathrm{t}}) = -1$. Bundles of higher topological charge can be constructed through the tensor product $E_{(k)} := E \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} E$. In fact $E_{(k)} = p_k(\mathcal{A})^{2^k}$, where $p_k := p \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} p \in M_{2^k}(\mathcal{A})$ is again a projector with $\operatorname{Tr} p_k = 1$, i.e. the bundles are complex line bundles, $\operatorname{rank}_{\mathbb{C}} E_{(k)} = 1$. Now

$$dp_k = \sum_{i=1}^{k} p \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} dp \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} p,$$

where dp appears once at the i-th position. Since $p(dp)p = 0$ and $\mathrm{Tr}\,(a \otimes_{\mathcal{A}} b) = \mathrm{Tr}\,a\,\mathrm{Tr}\,a$ we get for the Chern character of $E_{(k)}$

$$F_k = \mathrm{Tr}\,p_k(dp_k)(dp_k) = k\mathrm{Tr}\,p(dp)(dp), \tag{6}$$

i.e. simply k times the Chern character of charge one, and consequently for the associated topological charges

$$c_1(p_k) = \frac{1}{2\pi i}\int_{S^2} F_k = k. \tag{7}$$

The transposed projectors provide the negative Chern numbers $c_1(p_k^{\mathrm{t}}) = -k$.

4 Fuzzy Line Bundles

The fuzzy sphere [11], [12] is defined by the relations

$$[X_a, X_b] = i\kappa\epsilon_{abc}X_c, \quad \sum_{a=1}^{3}(X_a)^2 = 1, \tag{8}$$

where $\{X_a\}_{a=1}^3$ generate the irreducible spin j representation of $su(2)$ and κ is not a continuous parameter but restricted to

$$\kappa = \frac{1}{\sqrt{j(j+1)}}. \tag{9}$$

The associative $\mathbb{C}$-algebra generated by the X_a's is $M_N(\mathbb{C}) =: \mathcal{A}_N$, the algebra of $N \times N$-matrices with $N = 2j+1$. The differential calculus on $\mathcal{A}_N$ to make it the fuzzy sphere is derivation based: One chooses the three derivations defined by $e_a := (1/\kappa)\,\mathrm{ad}\,X_a$, satisfying $[e_a, e_b] = i\epsilon_{abc}e_c$, as analogue of the set of vector fields on the 2-sphere. One-forms df for $f \in \mathcal{A}_N$ are given through

$$df(u) = u(f), \quad \text{where } u = u^a e_a,$$

and $u^a \in \mathbb{C}$. Particularly $dX_a(u) = u(X_a)$. One can choose a basis θ^a of $\Omega^1(\mathcal{A}_N)$ completely determined by $\theta^a(e_b) = \delta^a_b$ such that $df = e_a(f)\theta^a$.

To proceed analogously to the commutative case we define $\mathbf{p} \in M_2(\mathcal{A}_N)$ through

$$\mathbf{p} := \alpha + \beta\,\sigma_a \otimes X_a, \tag{10}$$

with $\alpha, \beta \in \mathbb{R}$. Then $\mathbf{p}$ is a projector if and only if

$$\beta = \pm\frac{1}{\sqrt{4+\kappa^2}} \text{ and } \alpha = \frac{1+\beta\kappa}{2}. \tag{11}$$

Accordingly, with this choice for α and β, $\mathbf{p}$ defines a projective right $\mathcal{A}_N$-module $\mathbf{E} = \mathbf{p}\mathcal{A}_N^2$. Note that

$$\mathrm{Tr}\,\mathbf{p} = 1 + \beta\kappa = 1 \pm \frac{1}{N},$$

which is unequal 1 as long as $\kappa \neq 0$. We shall refer therefore to this noncommutative "line" bundles as fuzzy line bundles, their "fiber dimension" deviating slightly[1] from 1. The first component $\mathbf{F}$ of the Chern character of $\mathbf{E}$ is given by

$$\mathbf{F} = \mathrm{Tr}\,\mathbf{p}(d\mathbf{p})(d\mathbf{p}) = \mathrm{Tr}\,\beta^2(\alpha + \beta\,\sigma_a \otimes X_a)(\sigma_b \otimes dX_b)(\sigma_c \otimes dX_c).$$

A calculation very similar to the commutative case plus taking into account the commutator of the coordinates leads to

$$\mathbf{F} = \gamma_\pm(N)\frac{i\epsilon_{abc}}{4}X_a dX_b \wedge dX_c, \tag{12}$$

where

$$\gamma_\pm(N) = \frac{(1-\frac{1}{N^2})^{3/2}(N \pm (N^2-2))}{N^2-3}. \tag{13}$$

Now for $f \in \mathcal{A}_N$ and

$$\omega := \frac{\epsilon_{abc}X_a dX_b \wedge dX_c}{8\pi} \in \Omega^2(\mathcal{A}_N)$$

define the integral trough

$$\int^\star f\omega = \mathrm{Tr}_N f, \quad \text{where } \mathrm{Tr}_N(\cdot) = \frac{1}{N}\mathrm{Tr}\,(\cdot), \tag{14}$$

with $\int^\star \omega = 1$. The 2-form ω is our noncommutative volume form, in the commutative limit it converges to the normalized volume form on S^2. Consequently the first Chern number of the fuzzy line bundle $\mathbf{E}$ is given by

$$c_1(\mathbf{p}) = \frac{1}{2\pi i}\int^\star \mathbf{F},$$

and one finds

$$c_1(\mathbf{p}) = \gamma_\pm(N) \tag{15}$$

for the topological charge. In the commutative limit $\kappa \to 0$, i.e. $N \to \infty$ the topological charge is ±1.

Bundles of higher topological charge will be treated using a more general approach in a forthcoming paper.

---

[1] We prefer to think of the noncommutativity as $\kappa \ll 1$.

5 Acknowledgements

HG wants to thank the Center for Advanced Studies (ZHS) and the Institute for Theoretical Physics, Leipzig University, for the Leibniz–Professorship 2000/2001. Both authors want to thank A. Strohmaier for fruitful discussions and CWR thanks S. Kolb for helpful remarks. This work was partly supported by the "Deutsche Forschungsgemeinschaft" within the scope of the "Graduiertenkolleg Quantenfeldtheorie" of the University of Leipzig.

References

[1] H. Grosse, C. Klimcik and P. Presnajder, "Topologically Nontrivial Field Configurations in Noncommutative Geometry", Commun. Math. Phys. **178** (1996) 507.

[2] H. Grosse and P. Presnajder, "A Noncommutative Regularization of the Schwinger Model", Lett. Math. Phys. **46** (1998) 61.

[3] S. Baez, A. P. Balachandran, B. Idri and S. Vaidya, "Monopoles and Solitons in Fuzzy Physics", Commun. Math. Phys. **208** (2000) 787.

[4] A. P. Balachandran and S. Vaidya, "Instantons and Chiral Anomaly in Fuzzy Physics", Int. J. Mod. Phys. A **16** (2001) 17.

[5] P. Valtancoli, "Projectors for the Fuzzy Sphere", Preprint, hep-th/0101189.

[6] G. Landi, "Deconstructing Monopoles and Instantons", Rev. Math. Phys. **12** (2000) 1367.

[7] G. Landi, "Projective Modules of Finite Type and Monopoles over S(2)", J. Geom. Phys. **37** (2001) 47.

[8] G. Landi, "An Introduction to Noncommutative Spaces and their Geometries", Lecture Notes in Physics **m51**, Springer-Verlag, 1997.

[9] M. Karoubi and C. Leruste, "Algebraic Topology via Differential Geometry", London Mathematical Society Lecture Notes Series **99**, Cambridge University Press, 1989.

[10] A. Connes, "Noncommutative Geometry", Academic Press, 1994.

[11] J. Madore, "The Fuzzy Sphere", Class. Quant. Grav. **9** (1992) 69.

[12] J. Madore, "An Introduction to Noncommutative Differential Geometry and its Physical Applications", London Mathematical Society Lecture Note Series **257**, Cambridge University Press, 2nd Ed., 1999.